

Crystal structure of 2-(2-naphthyl)-4,6-dimethylpyrimidine, C₁₆H₁₄N₂

Li-Ming Qiang^I, Jun-Ling Si^{II}, Xin-Ming Dong^{III}, Xin-Qi Hao^{III}, Kui Lu^{*I} and Yu-Fen Zhao^{IV}


^I Henan Institute of Engineering, Department of Material and Chemical Engineering, Zhengzhou 470007, P. R. China

^{II} Zhengzhou University of Light Industry, School of Food and Biological Engineering, Zhengzhou 450002, P. R. China

^{III} Zhengzhou University, Department of Chemistry, Zhengzhou 450052, P. R. China

^{IV} Xiamen University, College of Chemistry and Chemical Engineering, Xiamen 361005, P. R. China

Received January 8, 2011, accepted and available on-line May 20, 2011; CCDC no. 1267/3361



Abstract

C₁₆H₁₄N₂, monoclinic, *P*12₁/*c*1 (no. 14), *a* = 8.387(1) Å, *b* = 17.453(3) Å, *c* = 8.694(1) Å, β = 93.092(2)°, *V* = 1270.8 Å³, *Z* = 4, *R*_{gt}(*F*) = 0.046, *wR*_{ref}(*F*²) = 0.129, *T* = 294 K.

Source of material

The title compound was obtained from the coupling reaction of 2-naphthaleneboronic acid and 2-iodo-4,6-dimethylpyrimidine as described in literature [1] and recrystallized from dichloromethane/petroleum ether solution at room temperature to give the desired crystals suitable for single crystal X-ray diffraction.

Experimental details

All the hydrogen atoms were positioned geometrically with *d*(C—H) = 0.93 Å, *U*_{iso}(H) = 1.2 *U*_{eq}(C) for the aromatic groups and *d*(C—H) = 0.96 Å, *U*_{iso}(H) = 1.5 *U*_{eq}(C) for the methyl groups. The H atoms of the methyl groups in the pyrimidine ring are disordered over two positions with occupation factors fixed at 0.5.

Discussion

Cyclometalated iridium complexes have attracted considerable attention in material research because of their outstanding performance in organic light-emitting diodes (OLED) [2,3]. In recent years, various types of cyclometalated Ir(III) complexes have been developed, such as homoleptic complexes, heteroleptic neutral complexes and cationic complexes [4–6]. In contrast to the most famous substituted phenylpyridine ligands, few examples of naphthalenylpyridine iridium complexes have been reported [7,8].

The pyrimidine ring and naphthalene ring in the crystal structure of the title compound are approximately coplanar with a dihedral

angle of 2.4°. There exist two types of intermolecular π–π stacking interactions. One π–π stacking interaction is between the pyrimidine rings (the inter-plane distance is 3.400 Å). The other π–π stacking interaction is between pyrimidine ring and naphthalene ring (the inter-plane distance is 3.748 Å). Both are attributed to construct the layer structure.

Table 1. Data collection and handling.

Crystal:	colorless block, size 0.03 × 0.31 × 0.42 mm
Wavelength:	Mo <i>K</i> _α radiation (0.71073 Å)
μ:	0.73 cm ^{−1}
Diffractometer, scan mode:	Bruker SMART CCD, φ/ω
2θ _{max} :	51°
<i>N</i> (<i>hkl</i>) _{measured} , <i>N</i> (<i>hkl</i>) _{unique} :	7463, 2367
Criterion for <i>I</i> _{obs} , <i>N</i> (<i>hkl</i>) _{gt} :	<i>I</i> _{obs} > 2 σ(<i>I</i> _{obs}), 1465
<i>N</i> (<i>param</i>) _{refined} :	164
Programs:	SHELXS-97, SHELXL-97, SHELXTL [9]

Table 2. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	Occ.	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> _{iso}
H(1)	4e		0.3477	0.5292	0.6776	0.058
H(3)	4e		0.0869	0.6172	0.5798	0.066
H(4)	4e		0.0189	0.7064	0.4241	0.069
H(6)	4e		0.2569	0.7642	0.3034	0.074
H(7)	4e		0.5251	0.7632	0.2707	0.081
H(8)	4e		0.6882	0.6728	0.3957	0.076
H(9)	4e		0.5846	0.5857	0.5581	0.066
H(13)	4e		0.1134	0.3600	0.9823	0.070
H(15A)	4e	0.50	0.3113	0.3528	0.9412	0.120
H(15B)	4e	0.50	0.1751	0.2913	0.9411	0.120
H(15C)	4e	0.50	0.2029	0.3466	1.0820	0.120
H(15D)	4e	0.50	0.1482	0.3076	1.0350	0.120
H(15E)	4e	0.50	0.2844	0.3692	1.0351	0.120
H(15F)	4e	0.50	0.2566	0.3138	0.8942	0.120
H(16A)	4e	0.50	0.3891	0.4981	0.8181	0.109
H(16B)	4e	0.50	0.3589	0.4632	0.9832	0.109
H(16C)	4e	0.50	0.3923	0.4091	0.8411	0.109
H(16D)	4e	0.50	0.3711	0.4154	0.9435	0.109
H(16E)	4e	0.50	0.4013	0.4504	0.7784	0.109
H(16F)	4e	0.50	0.3679	0.5045	0.9204	0.109

* Correspondence author (e-mail: lmq06@126.com)

Table 3. Atomic coordinates and displacement parameters (in Å²).

Atom	Site	<i>x</i>	<i>y</i>	<i>z</i>	<i>U</i> ₁₁	<i>U</i> ₂₂	<i>U</i> ₃₃	<i>U</i> ₁₂	<i>U</i> ₁₃	<i>U</i> ₂₃
C(1)	4e	0.2820(2)	0.5648(1)	0.6258(2)	0.045(1)	0.051(1)	0.049(1)	0.0058(8)	0.0011(9)	0.0031(9)
C(2)	4e	0.1210(2)	0.5624(1)	0.6445(2)	0.042(1)	0.051(1)	0.045(1)	0.0045(8)	0.0020(8)	0.0076(9)
C(3)	4e	0.0225(2)	0.6175(1)	0.5672(2)	0.042(1)	0.061(1)	0.062(1)	0.0081(9)	0.0025(9)	0.004(1)
C(4)	4e	0.0861(2)	0.6710(1)	0.4744(2)	0.052(1)	0.057(1)	0.063(1)	0.0140(9)	0.002(1)	0.001(1)
C(5)	4e	0.2516(2)	0.6742(1)	0.4527(2)	0.054(1)	0.046(1)	0.050(1)	0.0043(9)	0.0003(9)	0.0070(9)
C(6)	4e	0.3211(3)	0.7280(1)	0.3549(2)	0.070(1)	0.052(1)	0.062(1)	0.004(1)	0.003(1)	0.005(1)
C(7)	4e	0.4810(3)	0.7274(1)	0.3351(3)	0.073(2)	0.062(1)	0.068(2)	0.009(1)	0.010(1)	0.007(1)
C(8)	4e	0.5793(2)	0.6732(1)	0.4111(2)	0.052(1)	0.069(1)	0.070(1)	0.009(1)	0.009(1)	0.003(1)
C(9)	4e	0.5174(2)	0.6209(1)	0.5071(2)	0.047(1)	0.059(1)	0.059(1)	0.0001(9)	0.0013(9)	0.002(1)
C(10)	4e	0.3515(2)	0.6195(1)	0.5306(2)	0.047(1)	0.048(1)	0.046(1)	0.0014(8)	0.0020(9)	0.0068(9)
C(11)	4e	0.0512(2)	0.5032(1)	0.7431(2)	0.043(1)	0.054(1)	0.044(1)	0.0009(9)	0.0017(8)	0.0110(9)
C(12)	4e	0.0910(2)	0.3986(1)	0.8973(2)	0.054(1)	0.056(1)	0.056(1)	0.0023(9)	0.007(1)	0.003(1)
C(13)	4e	0.0706(2)	0.3975(1)	0.9207(2)	0.057(1)	0.061(1)	0.058(1)	0.010(1)	0.009(1)	0.003(1)
C(14)	4e	0.1669(2)	0.4524(1)	0.8516(2)	0.045(1)	0.069(1)	0.051(1)	0.008(1)	0.0056(9)	0.014(1)
C(15)	4e	0.2053(3)	0.3423(1)	0.9721(3)	0.070(2)	0.071(1)	0.100(2)	0.008(1)	0.011(1)	0.024(1)
C(16)	4e	0.3426(2)	0.4560(1)	0.8756(3)	0.045(1)	0.095(2)	0.078(2)	0.007(1)	0.010(1)	0.007(1)
N(1)	4e	0.1527(2)	0.45174(8)	0.8067(2)	0.0465(9)	0.0518(9)	0.056(1)	0.0018(7)	0.0069(8)	0.0002(8)
N(2)	4e	0.1066(2)	0.50631(9)	0.7609(2)	0.0422(9)	0.065(1)	0.051(1)	0.0008(7)	0.0038(7)	0.0089(8)

Acknowledgments. We thank the program for innovation of science and technology talents of Henan Province (grant no. 104200510022), the program for science and technology leaders of Zhengzhou City (grant no. 10LJRC174), the Doctor Foundation (grant no. D09004) and the program for innovative research team of Henan Institute of Engineering (grant no. 2009IRTHNIE05) for financial support.

References

- Wang, Z. Q.; Xu, C.; Dong, X. M.; Zhang, Y. P.; Hao, X. Q.; Gong, J. F.; Song, M. P.; Ji, B. M.: Synthesis, structure and properties of a novel iridium(III) pyrimidine complex. *Inorg. Chem. Commun.* **14** (2011) 316-319.
- Hsu, N. M.; Li, W. R.: Accelerated discovery of red-phosphorescent emitters through combinatorial organometallic synthesis and screening. *Angew. Chem., Int. Ed.* **45** (2006) 4138-4143.
- Chen, Z. Q.; Bian, Z. Q.; Huang, C. H.: Functional Ir(III) complexes and their applications. *Adv. Mater.* **22** (2010) 1534-1536.
- You, Y.; Park, S. Y.: Inter-ligand energy transfer and related emission change in the cyclometalated heteroleptic iridium complex: facile and efficient color tuning over the whole visible range by the ancillary ligand structure. *J. Am. Chem. Soc.* **127** (2005) 12438-12439.
- Huo, S. Q.; Deaton, J. C.; Rajeswaran, M.; Lenhart, W. C.: Highly efficient, selective, and general method for the preparation of meridional homo- and heteroleptic tris-cyclometalated iridium complexes. *Inorg. Chem.* **45** (2006) 3155-3157.
- Talarico, A. M.; Aiello, I.; Bellusci, A.; Crispini, A.; Ghedini, M.; Godbert, N.; Pugliese, T.; Szerb, E.: Highly luminescent bis-cyclometalated iridium(III) ethylenediamine complex: synthesis and correlation between the solid state polymorphism and the photophysical properties. *Dalton Trans.* (2010) 1709-1712.
- Yang, C. H.; Chen, C. H.; Sun, I. W.: Bathochromic effect of trifluoromethyl-substituted 2-naphthalen-1-yl-pyridine ligands in color tuning of iridium complexes. *Polyhedron* **25** (2006) 2407-2414.
- Zhen, H. Y.; Luo, C.; Yang, W.; Song, W. Y.; Du, B.; Jiang, J. X.; Jiang, C. Y.; Zhang, Y.; Cao, Y.: Electrophosphorescent chelating copolymers based on linkage isomers of naphthylpyridine-iridium complexes with fluorene. *Macromolecules* **39** (2006) 1693-1700.
- Sheldrick, G. M.: A short history of SHELX. *Acta Crystallogr.* **A64** (2008) 112-122.